

CO6: Introduction to Computational Neuroscience

<https://lnc2.dec.ens.fr/en/teams/mathematics-neural-circuits/co6-course>

Lecturer: Boris Gutkin
boris.gutkin@ens.fr
École Normale Supérieure
29 rue d'Ulm, 2nd floor

Exercise Sheet 4 — 10 April 2018

Please submit your solution in the next class (17 April 2018)

1. **Covariance and Correlation.** Assume that we have recorded two neurons in the two-alternative-forced choice task discussed in class. We denote the firing rate of neuron 1 in trial i as $r_{1,i}$ and the firing rate of neuron 2 as $r_{2,i}$. We furthermore denote the averaged firing rate of neuron 1 as \bar{r}_1 and of neuron 2 as \bar{r}_2 . Let us “center” the data by defining two data vectors

$$\mathbf{x} = \begin{pmatrix} r_{1,1} - \bar{r}_1 \\ r_{1,2} - \bar{r}_1 \\ r_{1,3} - \bar{r}_1 \\ \vdots \\ r_{1,N} - \bar{r}_1 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} r_{2,1} - \bar{r}_2 \\ r_{2,2} - \bar{r}_2 \\ r_{2,3} - \bar{r}_2 \\ \vdots \\ r_{2,N} - \bar{r}_2 \end{pmatrix}.$$

- (a) Show that the variance of the firing rates of the first neuron is

$$\text{Var}(r_1) = \frac{1}{N-1} \|\mathbf{x}\|^2.$$

- (b) Compute the cosine of the angle between \mathbf{x} and \mathbf{y} . What do you get?
(c) What are the maximum and minimum values that the correlation coefficient between r_1 and r_2 can take? Why?
(d) What do you think the term “centered” refers to?

2. **Bayes' theorem.** The theorem of Bayes summarizes all the knowledge we have about about the stimulus by observing the responses of a set of neurons, *independently* of the specific decoding rule. To get a better intuition about this theorem, we will look at the motion discrimination task again and compute the probability that the stimulus moved to the left (\leftarrow) or right (\rightarrow). For a stimulus $s = \{\leftarrow, \rightarrow\}$, and a firing rate response r of a single neuron, Bayes' theorem reads

$$p(s|r) = \frac{p(r|s)p(s)}{p(r)}.$$

Here, $p(r|s)$ is the probability that the firing rate is r if the stimulus was s . The respective distribution can be measured and we assume that it follows a Gaussian probability density

with mean μ_s and standard deviation σ ,

$$p(r|s) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(r - \mu_s)^2}{2\sigma^2}\right)$$

The relative frequency with which the stimuli (leftward or rightward motion, \leftarrow or \rightarrow) appear is denoted by $p(s)$, often called the *prior probability* or, for short, the *prior*. The distribution $p(r)$ denotes the probability of observing a response r , independent of any knowledge about the stimulus.

- How can you calculate $p(r)$? What shape does it have?
- The distribution $p(s|r)$ is often called the *posterior* probability or, for short, the *posterior*. Calculate the posterior for $s = \leftarrow$ and sketch it as a function of r , assuming a prior $p(\leftarrow) = p(\rightarrow) = 1/2$. Draw the posterior $p(\rightarrow | r)$ into the same plot.
- What happens if you change the prior? Investigate how the posterior changes if $p(\leftarrow)$ becomes much larger than $p(\rightarrow)$ and vice versa. Make a sketch similar to (b).
- Let us assume that we decide for leftward motion whenever $r > \frac{1}{2}(\mu_{\leftarrow} + \mu_{\rightarrow})$. Interpret this decision rule in the plots above. How well does this rule do depending on the prior? What do you lose when you move from the full posterior to a simple decision (decoding) rule?

3. Linear discriminant analysis (advanced): Let us redo the calculations for the case of N neurons. If we denote by \mathbf{r} the vector of firing rates, Bayes' theorem reads:

$$p(s|\mathbf{r}) = \frac{p(\mathbf{r}|s)p(s)}{p(\mathbf{r})} .$$

We assume that the distribution of firing rates again follows a Gaussian so that

$$p(\mathbf{r}|s) = \frac{1}{(2\pi)^{N/2} \sqrt{\det \mathbf{C}}} \exp\left(-\frac{1}{2}(\mathbf{r} - \boldsymbol{\mu}_s)^T \mathbf{C}^{-1}(\mathbf{r} - \boldsymbol{\mu}_s)\right)$$

where $\boldsymbol{\mu}_s$ denotes the mean of the density for stimulus $s = \{\leftarrow, \rightarrow\}$, and \mathbf{C} is the covariance matrix, assumed identical for both stimuli.

- Compute the log-likelihood ratio

$$l(\mathbf{r}) = \log \frac{p(\mathbf{r} | \leftarrow)}{p(\mathbf{r} | \rightarrow)} .$$

- Assume that $l(\mathbf{r}) = 0$ is the decision boundary, so that any firing rate vector \mathbf{r} giving a log-likelihood ratio larger than zero is classified as coming from the stimulus \leftarrow . Compute a formula for the decision boundary. What shape does this boundary have?
- Assume that $p(\leftarrow) = p(\rightarrow) = 1/2$. Assume we are analyzing two neurons with uncorrelated activities, so that the covariance matrix is

$$\mathbf{C} = \begin{pmatrix} \sigma_1^2 & 0 \\ 0 & \sigma_2^2 \end{pmatrix}$$

Sketch the decision boundary for this case.